Table 1.25 Age-Adjusted SEER Incidence Rates and Trends for the Top 15 Cancer Sites by Race/Ethnicity Females

| All Ra | ces | _ | White | | | Black | | | |
|-------------------------|--------------------------------|-------------------------------|----------------------------|--------------------------------|-------------------------------|-------------------------|--------------------------------|-------------------------------|--|
| | Rate ^b 2002-2006 | APC ^c 1997-2006 | | Rate ^b 2002-2006 | APC ^c 1997-2006 | | Rate ^b 2002-2006 | APC ^c 1997-2006 | |
| All Sites | 408.5 | -0.8* | All Sites | 420.5 | -0.8* | All Sites | 398.9 | -0.4* | |
| Breast | 123.8 | -1.8* | Breast | 127.8 | -1.9* | Breast | 117.7 | -0.6* | |
| Lung and Bronchus | 52.5 | -0.6* | Lung and Bronchus | 54.8 | -0.7* | Lung and Bronchus | 54.7 | 0.3 | |
| Colon and Rectum | 42.8 | -2.1* | Colon and Rectum | 42.1 | -2.3* | Colon and Rectum | 53.5 | -1.3* | |
| Corpus and Uterus, NOS | 23.3 | -0.8* | Corpus and Uterus, NOS | 24.2 | -1.0* | Corpus and Uterus, NOS | 20.3 | 1.3 | |
| Non-Hodgkin Lymphoma | 16.4 | 0.4 | Melanoma of the Skin | 18.7 | 2.1* | Pancreas | 14.6 | 0.2 | |
| Melanoma of the Skin | 15.8 | 1.6* | Non-Hodgkin Lymphoma | 17.2 | 0.5 | Non-Hodgkin Lymphoma | 12.2 | 0.7 | |
| Thyroid | 14.2 | 6.2* | Thyroid | 15.0 | 6.5* | Cervix Uteri | 10.4 | -5.2* | |
| Ovary ^g | 13.1 | -1.6* | Ovary ^g | 13.8 | -1.7* | Kidney and Renal Pelvis | 10.3 | 1.6 | |
| Pancreas | 10.4 | 0.6* | Pancreas | 10.2 | 0.8* | Ovary ^g | 10.1 | -1.0 | |
| Kidney and Renal Pelvis | 9.5 | 2.8* | Urinary Bladder | 9.9 | -0.6 | Myeloma | 10.0 | -0.4 | |
| Leukemia | 9.5 | -0.5 | Leukemia | 9.9 | -0.6 | Stomach | 9.0 | -3.1* | |
| Urinary Bladder | 9.3 | -0.6* | Kidney and Renal Pelvis | 9.9 | 2.9* | Thyroid | 8.4 | 6.8* | |
| Cervix Uteri | 8.2 | -3.3* | Cervix Uteri | 8.1 | -2.8* | Urinary Bladder | 7.9 | 0.3 | |
| Oral Cavity and Pharynx | 6.1 | -1.4* | Oral Cavity and Pharynx | 6.1 | -1.5* | Leukemia | 7.8 | -0.3 | |
| Stomach | 5.5 | -1.3* | Brain and ONS ^f | 5.9 | -0.7 | Oral Cavity and Pharynx | 5.8 | -1.1 | |

| Asian/Pacific | c Islander | | American Indian/Alaska Natived | | | Hispanic ^e | | |
|--------------------------|--------------------------------|-------------------------------|--------------------------------|--------------------------------|-------------------------------|--------------------------|--------------------------------|-------------------------------|
| | Rate ^b 2002-2006 | APC ^c 1997-2006 | | Rate ^b 2002-2006 | APC ^c 1997-2006 | | Rate ^b 2002-2006 | APC ^c 1997-2006 |
| All Sites | 287.5 | -0.5 | All Sites | 302.2 | 1.0 | All Sites | 312.5 | -0.6* |
| Breast | 89.5 | -0.9* | Breast | 74.4 | 1.4 | Breast | 88.3 | -0.6 |
| Colon and Rectum | 34.6 | -1.2 | Colon and Rectum | 41.2 | -0.5 | Colon and Rectum | 32.2 | -0.9* |
| Lung and Bronchus | 28.1 | 0.3 | Lung and Bronchus | 39.8 | 4.2 | Lung and Bronchus | 24.7 | -1.9* |
| Corpus and Uterus, NOS | 16.8 | 0.6 | Corpus and Uterus, NOS | 16.3 | 1.5 | Corpus and Uterus, NOS | 17.8 | 1.0 |
| Thyroid | 14.2 | 4.1* | Kidney and Renal Pelvis | 14.1 | - | Non-Hodgkin Lymphoma | 14.6 | 0.2 |
| Non-Hodgkin Lymphoma | 10.9 | -0.6 | Ovary ^g | 10.8 | -2.2 | Thyroid | 12.9 | 5.6* |
| Stomach | 10.0 | -3.3* | Non-Hodgkin Lymphoma | 10.0 | - | Cervix Uteri | 12.7 | -3.8* |
| Ovary ^g | 9.9 | -0.6 | Pancreas | 9.3 | - | Ovary ^g | 11.3 | -0.9 |
| Liver & IBD ^f | 8.3 | 0.8 | Thyroid | 8.5 | -0.2 | Pancreas | 10.2 | -0.8 |
| Pancreas | 8.2 | -0.2 | Stomach | 7.9 | - | Kidney and Renal Pelvis | 9.6 | 1.9* |
| Cervix Uteri | 7.6 | -4.6* | Cervix Uteri | 7.2 | 0.0 | Stomach | 9.3 | -1.0 |
| Leukemia | 6.1 | -0.1 | Liver & IBD ^f | 6.6 | - | Leukemia | 8.1 | -0.8 |
| Oral Cavity and Pharynx | 5.4 | -1.5 | Leukemia | 6.1 | _ | Liver & IBD ^f | 5.9 | 0.5 |
| Kidney and Renal Pelvis | 4.8 | 3.4* | Oral Cavity and Pharynx | 5.1 | - | Urinary Bladder | 5.3 | 0.4 |
| Urinary Bladder | 4.0 | -1.4 | Myeloma | 5.1 | - | Melanoma of the Skin | 4.7 | -1.7 |

- Top 15 cancer sites selected based on 2002-2006 age-adjusted rates for the race/ethnic group.
- Incidence data used in calculating the rates are from the 17 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry, Rural Georgia, California excluding SF/SJM/LA, Kentucky, Louisiana and New Jersey).
- Rates are age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- The APC is the Annual Percent Change over the time interval. Incidence data used in calculating the trends are from the 13 SEER areas (San Francisco, Connecticut, Detroit, Hawaii, Iowa, New Mexico, Seattle, Utah, Atlanta, San Jose-Monterey, Los Angeles, Alaska Native Registry and Rural Georgia).
- Trends are based on rates age-adjusted to the 2000 US Std Population (19 age groups Census P25-1130).
- d Rates for American Indian/Alaska Native are based on the CHSDA(Contract Health Service Delivery Area) counties.
- Hispanic is not mutually exclusive from whites, blacks, Asian/Pacific Islanders, and American Indians/Alaska Natives. Incidence data for Hispanics are based on NHIA and exclude cases from the Alaska Native Registry.
- IBD = Intrahepatic Bile Duct. ONS = Other Nervous System.
- Ovary excludes borderline cases or histologies 8442, 8451, 8462, 8472, and 8473.
- The APC is significantly different from zero (p<.05).

Statistic not shown. Rate based on less than 16 cases for the time interval. Trend based on less than 10 cases for at least one year within the time interval.